



Fumagri® OPP

Food safety begins
in your poultry sheds

THE TOOL FOR FIGHTING SALMONELLA, THE ALTERNATIVE TO FORMALDEHYDE


PRODUCT

- Bactericidal and fungicidal disinfectant with OrthoPhenylPhenol (OPP)
- Dispersion by ULTRADIFFUSION®
- Complies with biocide regulations, use PTO3: livestock premises disinfection

ADVANTAGES

- Demonstrated effectiveness against salmonella, aspergillus and other bacteria, and moulds
- Uniform diffusion of the disinfectant
- Safe to operators and animals: no humans or animals presence during treatment
- Convenient and fast application
- Non-corrosive

RESULTS




Required efficacy according to standard NFT 72281: 5 log for bacteria and 4 log for moulds




Fumagri® **OPP**

*Food safety begins
in your poultry sheds*

APPLICATION : Fumagri® OPP 1250 m³ (for treating volumes up to 1250 m³)


USE	<p>To prevent and eradicate fungal and bacterial contamination</p> <p>Optimal application conditions: Minimum temperature: 9°C Humidity 50-80%</p>	
------------	--	---



Type of shed	Number of Fumagri® OPP 1250m ³ tins
400m ²	1
800m ²	2
1000m ²	3
1200m ²	4 (3 for side access shed)
Laying premises	To suit shed volume

Remember to disinfect the silo with Fumagri® OPP 20g
2 days before feed loading,
Dry treatment, no humidity input.

Use PTO4: storage of livestock feed.



Also available for other applications in 20g, 40g, 120g, 200g, 400g pre-dosed tins.

Fumagri® OPP: biocidal use PTO3 and PTO4. Composition: OrthoPhenylPhenol, Xi classified, non-corrosive.
Take care when using biocides. Read label and product information before using.
Technical data sheet on request
design & layout: eretic brand consulting & design - photo credits: ©iStockphoto.com/barbara gibbons